An official website of the United States government Here's how you know ▾

Springer Journal of Environmental Health Science & Engineering

J Environ Health Sci Eng. 2015; 13: 34.
Published online 2015 Apr 21. doi: 10.1186/s40201-015-0192-2
PMCID: PMC4416385
PMID: 25937951

Use of mobile phone during pregnancy and risk of spontaneous abortion

Fatemeh Shamsi Mahmoodabadi, Saeedeh Ziaei,* Mohammad Firoozabadi, and Anushiravn Kazemnejad
› Author information › Article notes › Copyright and License information Disclaimer

Abstract

Background

Exposure to electromagnetic fields of cell phones increasingly occurs, but the potential influence on spontaneous abortion has not been thoroughly investigated.

Methods

We carried out a prospective cohort study on 1300 pregnant women in the first trimester of pregnancy. Mobile phone users were considered as exposed subjects and the others as unexposed. Women were followed until the pregnancy ended in spontaneous abortion (case group) or normal delivery (control group). Data were collected through a questionnaire and analyzed by logistic regression and survival analysis.

Results

All the data pertaining to mobile phones were different between the two groups except the use of hands-free devices ($p < 0.001$).

Conclusion

Our result suggests that use of mobile phones can be related to the early spontaneous abortions.

Keywords: Abortion, Electromagnetic fields, Mobile phones, Pregnancy

Introduction

Widespread concerns have been raised about possible effects of electromagnetic field (EMF) from personal and mobile telecommunication [1-3]. Much of the concern arises because new technologies are introduced without adequate previous investigation into their potential harm and nature or discussion of the debate within the scientific community about possible health consequences. Indeed, mobile phone use has increased considerably along with reducing its costs, and developing countries are establishing mobile telecommunications rather than the more expensive fixed-line systems. Thus, if there is an impact on health, it is likely to be greater and to affect everyone in the world. The major focus of research has been on radio-frequency (RF) radiation, which is mainly generated by antennas, while some scientists are concerned about the possible impact of exposure to extremely low frequency electromagnetic field (ELF-EMF) fields generated by supply currents in the power [4]. The focus of the present study on electromagnetic radiation is the front-end amplifier. Consequently, the corresponding ELF-EMF has a spectrum similar to the pulse structure of RF signals.

The public discussion of the health hazards of EMF exposure has focused on the possible association with cancer, cardiovascular and immune systems. Less attention has been paid to the evaluation of evaluating the health hazards of EMF exposure on reproductive health such as spontaneous abortion [5-9].

The health effect of EMF on pregnancy has remained controversial despite efforts to reach consensus [10]. The main challenge is studying EMF area: 1) accurately assessing during the relevant time period, and 2) identifying susceptible population.

Therefore a case-control study has been designed o determine the effects of EMF exposure to mobile phones immediately after spontaneous abortions in selected patients.

Methods

For the evaluation of the effects of EMF exposure to mobile phones on spontaneous abortions, a case-control study was chosen. Due to the lack of knowledge about these effects, a pilot study on 100 participants was carried out. Then, based on the primary data, with a significance level of $\alpha = 0.05$ and power of $1-\beta = 0.80$,a sample size of 600 cases and 600 controls was calculated. In the current case-control study, 2592 women who had an unexplained spontaneous abortion at < 14 weeks of gestation (as early abortion) immediately after abortion(during first 48 hours) and 300 pregnant women at 14 weeks gestation were enrolled. Gestational age was calculated based on the last menstrual period that was confirmed by an ultrasonographic examination. The subjects were recruited from 10 hospitals in Tehran.

Pregnancy situations in the pregnant women in the control group were matched with maternal and paternal age, gravidity, pre-pregnancy BMI, family relationship, duration from last delivery, educational level, occupation and history of abortions or preterm labor with the women in the case group.

The inclusion criteria were as follows:

1. singleton pregnancy; 2) 18–35 years of age; 3) spontaneous pregnancy without the use of assisted reproductive technology

The exclusion criteria were as follows:

1. chronic diseases (diabetes, hypertension,thyroid diseases, immune system diseases and cardiovascular diseases); 2) any genetic disorders (in the women, their husbands,and their first-degree relatives);3) vaginal bleeding in the first trimester of pregnancy in the control group; 4) history of birth defects in previous pregnancies; and 5) cigarette smoking, alcohol or any drug consumptions.

The women completed the data collection form, which was used to collect data about sociodemographic and obstetric characteristics, medical and reproductive histories, and lifestyles. The second data collection form which was completed by trained interviewer was designed to obtain information about the use of cell phones during pregnancy. For the consideration of cell phones radiation, we asked the women about the age at first initiation of the cell phone,when was not in use (in handbag, clothing pocket, or in60-70 cm away from the body), use of hands-free equipment, use of flowers for other applications (in send messages, to listen to music, and to play games), the specific absorption rate (SAR) reported by the manufacturer and the average of the effective SAR (average duration of calling time per day × SAR). SAR is a measure which electromagnetic energy is absorbed by body tissues. This parameter shows the estimation of total EMF energy. The effective SAR can show the exposure of ELF-estimation from mobile phone indirectly. We investigated the data about the SAR of each mobile phone model from SARfer site.

For precise evaluation of the calling time, the records of mobile phone bills were considered. The women were asked about the time spent for telecommunications in the specific section of were reviews of cell phones were also taken into consideration.

The procedure was approved by the Ethics Committee of Tarbiat Modares University. All of the women participated voluntarily and provided written informed consent.

Statistical analysis

Demographic and obstetric characteristics, as well as the parameters which reflect the use of cell phones were compared between the two groups using t-tests and chi-square tests. A logistic regression model was used to assess the association between spontaneous abortions as a binary dependent factor and effective SAR, maternal and paternal age, history of abortions and family relationship as independent factors. Analyses were carried out with statistical package state software(SPSS) ULS Fvalue v10.05 was considered statistically significant. All confidence intervals were calculated at the 95% level.

Results

The distribution of demographic and potential confounders for spontaneous abortions is presented in Table 1. As shown, there were no significant differences in maternal and paternal ages, pre-pregnancy BMI, occupation and educational status, socioeconomic level, family relationship, duration from last delivery and history of previous abortions or preterm labors between the two groups.

Table 1

The comparison of the maternal and paternal characteristics between the two groups

Parameters	Case groups:n (2326)	Control groups:n (346)	P
Maternal age(years)* Mean ± SD	27.81 ± 5.20	27.34 ± 4.1	0.2
Paternal age(years)* Mean ± SD	32.62 ± 5.46	31.98 ± 5.49	0.1
Family relationship* N(%)	2322(1.0)	772(5.1)	0.1
Pre-pregnancy BMI*	25.14 ± 4.98	24.29 ± 5.24	0.09
History of abortion** N(%)	47(20.0)	36(14.6)	0.2
History of preterm labor** N(%)	5(1.5)	2(0.8)	0.6
Duration from last delivery(years)* Mean ± SD	6.61 ± 3.93	6.57 ± 5.46	0.9
Occupation N(%)**			0.2
Housewife	195(86.4)	225(95.5)	
Employee	31(13.6)	21(8.5)	
Educational status N(%)**			0.09
Primary school	88(38.9)	98(39.8)	
Secondary and high school	99(43.4)	108(46.6)	
University	44(19.5)	48(19.6)	

Open in a separate window

* Student test **Chi square test.

As shown in Table 2, all of the data pertaining to the use of cell phones such as average calling time per day, the location of the cell phone when not in use, use of phones for other applications, the specific absorption rate (SAR) and the average of the effective SAR), except the use of hands free devices were different between the two groups ($p < 0.001$). Logistic regression analysis revealed a significant association between the effective SAR with the risk of spontaneous abortion after adjustment for maternal age, paternal age, history of abortion and family relationships (OR: 1.11,p < 0.001) (Table 3).

Table 2

The comparison of cell phone use during pregnancy between the two groups

Parameters	Case groups:n (236)	Control groups:n (346)	P(2-tailed)
Calling time per day* (minutes) Mean ± SD	9.31 ± 11.22	5.14 ± 8.39	<0.001
Use of hands-free** n(%)	19(8.6)	15(4.5)	0.09
location of phones when not in use** n(%)			<0.001
on tables	86(38.1)	77(31.3)	
in handbags	101(44.7)	153(62.2)	
in pocket	39(17.3)	16(6.5)	
use of phone for other applications **(%)	95(39.8)	44(17.9)	<0.001
Effective SAR* Mean ± SD	7.02 ± 13.92	2.28 ± 3.20	<0.001

Open in a separate window

* Student test, ** Chi square test or Fisher's exact test,SAR: Specific Absorption Rate.

Table 3

Association of spontaneous abortions with the effective SAR(Specific Absorption Rate) *

	OR(95%CI)	B	SE	P
Effective SAR	1.11(1.07-1.16)	0.106	0.21	<0.001

Open in a separate window

*Adjusted for maternal age, paternal age, history of abortion and the spouse relatives.

Discussion

Exposure to EMF is increasingly common, but the potential influence on health has not been thoroughly determined, especially in pregnant women. Electromagnetic fields may produce chromosome and even free radicals, which can make a susceptible population prone to congenital malformations and tissue and cell damages [11]. Long-term exposure to electromagnetic fields may be linked to even higher levels of oxidative stress, with the aforementioned corresponding consequences.

There is limited information on the association between EMF exposure during pregnancy and reproductive outcomes [10-18]. Some studies have reported increased risk of spontaneous abortions and congenital malformations, although these results were derived from poorly designed studies.

This study demonstrated an increased risk of spontaneous abortions associated with EMF exposure and confirmed the results of other researches [10,12]. Lee conducted a nested case-control study to evaluate the association of residential and occupational magnetic fields with selected spontaneous abortions [12]. He reported personal magnetic field maximum exposure is associated with a higher risk of spontaneous abortion, but not with residential magnetic field exposure, specifically the magnetic field measurements were collected in the second trimester and the occurrence of the spontaneous abortions. Perhaps the behavior and magnetic fields of women who had spontaneous abortions were different from the pregnant controls. But our study had this strength that, the data were obtained shortly after the spontaneous abortions occurred in the case group,and in the early second trimester in the control group. This led to a better evaluation of the environmental conditions and the lifestyles of the participants

Though, the present findings are inconsistent with the results of some other works [13-18]. Some studies do not support an association between electric and magnetic fields and reproductive damages. However many of them had investigated the effects of EMF either on animals or in vitro conditions [13-15]. Also, lack of knowledge about the dose for a strong association between EMF and adverse reproductive outcomes may be primarily the result of having little information data due to the lack of epidemiological data in this area.

One potential limitation need to be kept in mind when one interprets the results of the current study. The study did not assess all the unknown risk factors for spontaneous abortions, such as balanced chromosomal abnormalities in one parent. Also, the data about unknown spontaneous abortion at very early stages were not collected. Other limitation of this study is its case control nature, and this can imply that caution should be taken in causal interpretations of the findings. The third limitation of this study is that cell phones may not be the only source of EMF. Since many parameters which can affect the fetal outcome, and the epidemiologic studies were not matched between the two groups, it is suspected that probably the effects that observed in this study are due to the use of cell phones.

Conclusion

Despite the lack of a clear understanding of the underlying mechanisms, the present result suggests that the use of cell phones may be related to spontaneous abortions, further study is warranted.

Acknowledgements

This study was carried out with the kind cooperation of the participating patients. Furthermore, this study was funded by Tarbiat Modares University, Tehran, Iran.

Footnotes

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

FSM carried out design and data collection of the study, SZ carried out design of the study and prepared the article, MF participated in design of the study, AK advised the statistical design of the study. All authors read and approved the final manuscript.

Correspondence Information

Fatemeh Shamsi Mahmoodabadi, Email: shamsi1@modares.ac.ir;

Saeedeh Ziaei, Email: ziaei_s@modares.ac.ir;

Mohammad Firoozabadi, Email: mozerm@modares.ac.ir;

Anushiravn Kazemnejad, Email: akk6@yahoo.com.

References

1. Repacholi MH. *Health risks from the use of mobile phone. Occupational and Environmental Health*. 1221 Geneva 27,Switzerland: World Health Organization, 2001. [Google Scholar]

2. Divan HA, Kheifets L, Obel C, Olsen J. Prenatal and postnatal exposure to cell phone use and behavioral problems in children. *Epidemiology*. 2008;19(1):1–7. doi: 10.1097/EDE.0b013e318157e457. [PubMed] [CrossRef] [Google Scholar]

3. Tsoutsman A. Human exposure by mobile phones in enclosed area. *Bioelectromagnetics*. 2003;24(3):63–5. doi: 10.1002/bem.10087. [PubMed] [CrossRef] [Google Scholar]

4. Wiley J, Sanderson W, Hansson MK. Subjective symptoms among mobile phone users-A consequence of absorption of radiofrequency fields? *Bioelectromagnetics*. 2003;24(7):132–9. doi: 10.1002/bem.10101. [PubMed] [CrossRef] [Google Scholar]

5. Robert E. Intracavitary effects of electromagnetic fields(low frequency, and frequency RF and microwave).review of epidemiologic studies. *Toxicology*. 1999;139:251–6. doi: 10.1016/S0300-483X(99)00029-4+292–[ADS: TRAA14:A19262] [PubMed] [CrossRef] [Google Scholar]

6. Berrada N, Tellaoueti E, Rasse V, Martelli A, Tavirat M, Mattioli M, et al. Extremely low frequency electromagnetic field exposure affects fertilization estimate in swine animal model. *Theriogenology*. 2010;73(7):1293–305. doi: 10.1016/j.theriogenology.2009.12.010. [PubMed] [CrossRef] [Google Scholar]

7. Juutilainen J. Effects of low-level radio frequency(RF) fields in 50HZon lymnaja on human cardiovascular, reproduction, immune, and other systems:A review of the recent literature. *Int J Hyg Environ Health*. 2010;213(1):2–3. 29. doi: 10.1016/j.ijheh.2007.05.001. [PubMed] [CrossRef] [Google Scholar]

8. Erdogan F. Electromagnetic pollution in the cities: The effects on the learning environment. *Essays Educ*. 2007;22:78–96. [Google Scholar]

9. Aldhous A, Castle E, Green A, Liser M, Sastre D, Swerdlow A. Review of the epidemiologic literature on EMF and health. *Environmental Health Perspect*. 2001;109(9):911–33. doi: 10.2307/3454651. [PMC free article] [PubMed] [CrossRef] [Google Scholar] [Ref list]

10. Li D-K, Odouli R, Wi S, Janevic T, Golditch I, Bracken TD, et al. A population-based prospective cohort study of personal exposure to magnetic fields during pregnancy and the risk of miscarriage. *Epidemiology*. 2002;13:9–20. doi: 10.1097/00001648-200201000-00004. [PubMed] [CrossRef] [Google Scholar]

11. Lee GM, Neutra RR, Hristova L, Yost M, Hiatt RA. A nested case-control study of residential and personal magnetic field measures and miscarriages. *Epidemiology*. 2002;13(21–31. doi: 10.1097/00001648-200201000-00005. [PubMed] [CrossRef] [Google Scholar]

12. Savitz DA. Magnetic fields and miscarriage. *Epidemiology*. 2002;13(1):1–3. doi: 10.1097/00001648-200201000-00001. [PubMed] [CrossRef] [Google Scholar]

13. Khaki AA, Khaki A, Garacharloo S, Khorshidi T, Tajadini M, Nadimi N. Pre and post natal exposure of 50 HZ electromagnetic fields on prostate gland of rats an electron microscopy study. *Iran J Reproductive Med*. 2006;4(2):77–82. [Google Scholar]

14. Sun W, Tan Q, Pan Y, Yu C, Sun X, Chang W. Effects of 50-Hz magnetic field exposure on hormonal secretion and chromosomal gene expression in human first-trimester villous trophoblasts in vitro. *Bioelectromagnetics*. 2010;31(7):561–72. doi: 10.1002/bem.20596. [PubMed] [CrossRef] [Google Scholar]

15. Hosseinnia H, Sasanianmoore Y, Kentsalamon K, Lotfinia J, Ivanlalaee J. Effects of low-frequency magnetic fields on implantation in rats. Reproductive Toxicology. 2000; 33(1):49–55. [PubMed]

16. Parsons P, Chalfoun F, Chatmoud L, Cipriani S, Ricci E, Chamery V, et al. Exposure to video display terminals and risk of small for- gestational age birth. *J Environ Health*. 2010;72(1):26–7. [PubMed] [Google Scholar]

17. Marcus M, McGeehin J, Golden A, Landrigan P. Video display terminals and miscarriage. *J Am Med Womens Assoc*. 2000;45(2):46–8. [PubMed] [Google Scholar]

18. Parazzini F, Chiaffarino F, Chatmoud L, Cipriani S, Ricci E, Chamery V, et al. Exposure to video display terminals and risk of small for- gestational age birth. *J Environ Health*. 2010;72(1):26–7. [PubMed] [Google Scholar]

Articles from Journal of Environmental Health Science and Engineering are provided here courtesy of Springer

FOLLOW NCBI

Connect with NLM
National Library of Medicine
8600 Rockville Pike
Bethesda, MD, 20894
NLM | NIH | HHS | USA.gov
Web Policies
FOIA
HHS Vulnerability Disclosure
Help
Accessibility
Careers